

A PARTIAL CAYLEY TRANSFORM OF SIEGEL-JACOBI DISK

JAE-HYUN YANG

ABSTRACT. Let $\mathbb{H}_g$ and $\mathbb{D}_g$ be the Siegel upper half plane and the generalized unit disk of degree g respectively. Let $\mathbb{C}^{(h,g)}$ be the Euclidean space of all $h \times g$ complex matrices. We present a partial Cayley transform of the Siegel-Jacobi disk $\mathbb{D}_g \times \mathbb{C}^{(h,g)}$ onto the Siegel-Jacobi space $\mathbb{H}_g \times \mathbb{C}^{(h,g)}$ which gives a partial bounded realization of $\mathbb{H}_g \times \mathbb{C}^{(h,g)}$ by $\mathbb{D}_g \times \mathbb{C}^{(h,g)}$. We prove that the natural actions of the Jacobi group on $\mathbb{D}_g \times \mathbb{C}^{(h,g)}$ and $\mathbb{H}_g \times \mathbb{C}^{(h,g)}$ are compatible via a partial Cayley transform. A partial Cayley transform plays an important role in computing differential operators on the Siegel-Jacobi disk $\mathbb{D}_g \times \mathbb{C}^{(h,g)}$ invariant under the natural action of the Jacobi group on $\mathbb{D}_g \times \mathbb{C}^{(h,g)}$ explicitly.

1. Introduction

For a given fixed positive integer g , we let

$$\mathbb{H}_g = \left\{ \Omega \in \mathbb{C}^{(g,g)} \mid \Omega = {}^t\Omega, \quad \text{Im } \Omega > 0 \right\}$$

be the Siegel upper half plane of degree g and let

$$Sp(g, \mathbb{R}) = \left\{ M \in \mathbb{R}^{(2g,2g)} \mid {}^tM J_g M = J_g \right\}$$

be the symplectic group of degree g , where $F^{(k,l)}$ denotes the set of all $k \times l$ matrices with entries in a commutative ring F for two positive integers k and l , tM denotes the transpose matrix of a matrix M and

$$J_g = \begin{pmatrix} 0 & I_g \\ -I_g & 0 \end{pmatrix}.$$

We see that $Sp(g, \mathbb{R})$ acts on $\mathbb{H}_g$ transitively by

$$(1.1) \quad M \cdot \Omega = (A\Omega + B)(C\Omega + D)^{-1},$$

where $M = \begin{pmatrix} A & B \\ C & D \end{pmatrix} \in Sp(g, \mathbb{R})$ and $\Omega \in \mathbb{H}_g$.

Let

$$\mathbb{D}_g = \left\{ W \in \mathbb{C}^{(g,g)} \mid W = {}^tW, \quad I_g - W\overline{W} > 0 \right\}$$

be the generalized unit disk of degree g . The Cayley transform $\Phi: \mathbb{D}_g \longrightarrow \mathbb{H}_g$ defined by

$$(1.2) \quad \Phi(W) = i(I_g + W)(I_g - W)^{-1}, \quad W \in \mathbb{D}_g$$

is a biholomorphic mapping of $\mathbb{D}_g$ onto $\mathbb{H}_g$ which gives the bounded realization of $\mathbb{H}_g$ by $\mathbb{D}_g$ (cf. [6, pp.281-283]). And the action (2.8) of the symplectic group on $\mathbb{D}_g$ is compatible with the action (1.1) via the Cayley transform Φ .

2000 Mathematics Subject Classification: Primary 32Hxx, 32M10; Secondary 11F50.

Keywords and phrases: Partial Cayley transform, Siegel-Jacobi space, Siegel-Jacobi disk, Harish-Chandra decomposition, automorphic factors, Jacobi forms.

For two positive integers g and h , we consider the Heisenberg group

$$H_{\mathbb{R}}^{(g,h)} = \left\{ (\lambda, \mu; \kappa) \mid \lambda, \mu \in \mathbb{R}^{(h,g)}, \kappa \in \mathbb{R}^{(h,h)}, \kappa + \mu^t \lambda \text{ symmetric} \right\}$$

endowed with the following multiplication law

$$(\lambda, \mu; \kappa) \circ (\lambda', \mu'; \kappa') = (\lambda + \lambda', \mu + \mu'; \kappa + \kappa' + \lambda^t \mu' - \mu^t \lambda').$$

The Jacobi group G^J is defined as the semidirect product of $Sp(g, \mathbb{R})$ and $H_{\mathbb{R}}^{(g,h)}$

$$G^J = Sp(g, \mathbb{R}) \ltimes H_{\mathbb{R}}^{(g,h)}$$

endowed with the following multiplication law

$$(M, (\lambda, \mu; \kappa)) \cdot (M', (\lambda', \mu'; \kappa')) = (MM', (\tilde{\lambda} + \lambda', \tilde{\mu} + \mu'; \kappa + \kappa' + \tilde{\lambda}^t \mu' - \tilde{\mu}^t \lambda'))$$

with $M, M' \in Sp(g, \mathbb{R})$, $(\lambda, \mu; \kappa), (\lambda', \mu'; \kappa') \in H_{\mathbb{R}}^{(g,h)}$ and $(\tilde{\lambda}, \tilde{\mu}) = (\lambda, \mu)M'$. Then G^J acts on $\mathbb{H}_g \times \mathbb{C}^{(h,g)}$ transitively by

$$(1.3) \quad (M, (\lambda, \mu; \kappa)) \cdot (\Omega, Z) = (M \cdot \Omega, (Z + \lambda\Omega + \mu)(C\Omega + D)^{-1}),$$

where $M = \begin{pmatrix} A & B \\ C & D \end{pmatrix} \in Sp(g, \mathbb{R})$, $(\lambda, \mu; \kappa) \in H_{\mathbb{R}}^{(g,h)}$ and $(\Omega, Z) \in \mathbb{H}_g \times \mathbb{C}^{(h,g)}$. In [9, p. 1331], the author presented the natural construction of the action (1.3).

We mention that studying the Siegel-Jacobi space or the Siegel-Jacobi disk associated with the Jacobi group is useful to the study of the universal family of polarized abelian varieties (cf. [10], [12]). The aim of this paper is to present a partial Cayley transform of the Siegel-Jacobi disk $\mathbb{D}_g \times \mathbb{C}^{(h,g)}$ onto the Siegel-Jacobi space $\mathbb{H}_g \times \mathbb{C}^{(h,g)}$ which gives a partially bounded realization of $\mathbb{H}_g \times \mathbb{C}^{(h,g)}$ by $\mathbb{D}_g \times \mathbb{C}^{(h,g)}$ and to prove that the natural actions of the Jacobi group on $\mathbb{D}_g \times \mathbb{C}^{(h,g)}$ and $\mathbb{H}_g \times \mathbb{C}^{(h,g)}$ are compatible via a partial Cayley transform. The main reason that we study a partial Cayley transform is that this transform is usefully applied to computing differential operators on the Siegel-Jacobi disk $\mathbb{D}_g \times \mathbb{C}^{(h,g)}$ invariant under the action (3.5) of the Jacobi group G_*^J (cf. (3.2)) explicitly.

This paper is organized as follows. In Section 2, we review the Cayley transform of the generalized unit disk $\mathbb{D}_g$ onto the Siegel upper half plane $\mathbb{H}_g$ which gives a bounded realization of $\mathbb{H}_g$ by $\mathbb{D}_g$. In Section 3, we construct a partial Cayley transform of the Siegel-Jacobi disk $\mathbb{D}_g \times \mathbb{C}^{(h,g)}$ onto the Siegel-Jacobi space $\mathbb{H}_g \times \mathbb{C}^{(h,g)}$ which gives a partially bounded realization of $\mathbb{H}_g \times \mathbb{C}^{(h,g)}$ by $\mathbb{D}_g \times \mathbb{C}^{(h,g)}$ (cf. (3.6)). We prove that the action (1.3) of the Jacobi group G^J is compatible with the action (3.5) of the Jacobi group G_*^J through a partial Cayley transform (cf. Theorem 3.1). In the final section, we present the canonical automorphic factors of the Jacobi group G_*^J .

NOTATIONS: We denote by $\mathbb{R}$ and $\mathbb{C}$ the field of real numbers, and the field of complex numbers respectively. For a square matrix $A \in F^{(k,k)}$ of degree k , $\sigma(A)$ denotes the trace of A . For $\Omega \in \mathbb{H}_g$, $\text{Re } \Omega$ (*resp.* $\text{Im } \Omega$) denotes the real (*resp.* imaginary) part of Ω . For a matrix $A \in F^{(k,k)}$ and $B \in F^{(k,l)}$, we write $A[B] = {}^tBAB$. I_n denotes the identity matrix of degree n .

2. The Cayley Transform

Let

$$(2.1) \quad T = \frac{1}{\sqrt{2}} \begin{pmatrix} I_g & I_g \\ iI_g & -iI_g \end{pmatrix}$$

be the $2g \times 2g$ matrix represented by Φ . Then

$$(2.2) \quad T^{-1}Sp(g, \mathbb{R})T = \left\{ \begin{pmatrix} P & Q \\ \overline{Q} & \overline{P} \end{pmatrix} \mid {}^tP\overline{P} - {}^t\overline{Q}Q = I_g, {}^tP\overline{Q} = {}^t\overline{Q}P \right\}.$$

Indeed, if $M = \begin{pmatrix} A & B \\ C & D \end{pmatrix} \in Sp(g, \mathbb{R})$, then

$$(2.3) \quad T^{-1}MT = \begin{pmatrix} P & Q \\ \overline{Q} & \overline{P} \end{pmatrix},$$

where

$$(2.4) \quad P = \frac{1}{2} \left\{ (A + D) + i(B - C) \right\}$$

and

$$(2.5) \quad Q = \frac{1}{2} \left\{ (A - D) - i(B + C) \right\}.$$

For brevity, we set

$$G_* = T^{-1}Sp(g, \mathbb{R})T.$$

Then G_* is a subgroup of $SU(g, g)$, where

$$SU(g, g) = \left\{ h \in \mathbb{C}^{(g, g)} \mid {}^thI_{g, g}\overline{h} = I_{g, g} \right\}, \quad I_{g, g} = \begin{pmatrix} I_g & 0 \\ 0 & -I_g \end{pmatrix}.$$

In the case $g = 1$, we observe that

$$T^{-1}Sp(1, \mathbb{R})T = T^{-1}SL_2(\mathbb{R})T = SU(1, 1).$$

If $g > 1$, then G_* is a *proper* subgroup of $SU(g, g)$. In fact, since ${}^tTJ_gT = -iJ_g$, we get

$$(2.6) \quad G_* = \left\{ h \in SU(g, g) \mid {}^thJ_g h = J_g \right\} = SU(g, g) \cap Sp(g, \mathbb{C}),$$

where

$$Sp(g, \mathbb{C}) = \left\{ \alpha \in \mathbb{C}^{(2g, 2g)} \mid {}^t\alpha J_g \alpha = J_g \right\}.$$

Let

$$P^+ = \left\{ \begin{pmatrix} I_g & Z \\ 0 & I_g \end{pmatrix} \mid Z = {}^tZ \in \mathbb{C}^{(g, g)} \right\}$$

be the P^+ -part of the complexification of $G_* \subset SU(g, g)$. We note that the Harish-Chandra decomposition of an element $\begin{pmatrix} P & Q \\ \overline{Q} & \overline{P} \end{pmatrix}$ in G_* is

$$\begin{pmatrix} P & Q \\ \overline{Q} & \overline{P} \end{pmatrix} = \begin{pmatrix} I_g & Q\overline{P}^{-1} \\ 0 & I_g \end{pmatrix} \begin{pmatrix} P - Q\overline{P}^{-1}\overline{Q} & 0 \\ 0 & \overline{P} \end{pmatrix} \begin{pmatrix} I_g & 0 \\ \overline{P}^{-1}\overline{Q} & I_g \end{pmatrix}.$$

For more detail, we refer to [2, p. 155]. Thus the P^+ -component of the following element

$$\begin{pmatrix} P & Q \\ \overline{Q} & \overline{P} \end{pmatrix} \cdot \begin{pmatrix} I_g & W \\ 0 & I_g \end{pmatrix}, \quad W \in \mathbb{D}_g$$

of the complexification of G_*^J is given by

$$(2.7) \quad \left(\begin{pmatrix} I_g & (PW + Q)(\overline{Q}W + \overline{P})^{-1} \\ 0 & I_g \end{pmatrix} \right).$$

We note that $Q\overline{P}^{-1} \in \mathbb{D}_g$. We get the Harish-Chandra embedding of $\mathbb{D}_g$ into P^+ (cf. [2, p. 155] or [5, pp. 58-59]). Therefore we see that G_* acts on $\mathbb{D}_g$ transitively by

$$(2.8) \quad \begin{pmatrix} P & Q \\ \overline{Q} & \overline{P} \end{pmatrix} \cdot W = (PW + Q)(\overline{Q}W + \overline{P})^{-1}, \quad \begin{pmatrix} P & Q \\ \overline{Q} & \overline{P} \end{pmatrix} \in G_*, \quad W \in \mathbb{D}_g.$$

The isotropy subgroup at the origin o is given by

$$K = \left\{ \begin{pmatrix} P & 0 \\ 0 & \overline{P} \end{pmatrix} \mid P \in U(g) \right\}.$$

Thus G_*/K is biholomorphic to $\mathbb{D}_g$. It is known that the action (1.1) is compatible with the action (2.8) via the Cayley transform Φ (cf. (1.2)). In other words, if $M \in Sp(g, \mathbb{R})$ and $W \in \mathbb{D}_g$, then

$$(2.9) \quad M \cdot \Phi(W) = \Phi(M_* \cdot W),$$

where $M_* = T^{-1}MT \in G_*$. For a proof of Formula (2.9), we refer to the proof of Theorem 3.1.

For $\Omega = (\omega_{ij}) \in \mathbb{H}_g$, we write $\Omega = X + iY$ with $X = (x_{ij})$, $Y = (y_{ij})$ real and $d\Omega = (d\omega_{ij})$. We also put

$$\frac{\partial}{\partial \Omega} = \left(\frac{1 + \delta_{ij}}{2} \frac{\partial}{\partial \omega_{ij}} \right) \quad \text{and} \quad \frac{\partial}{\partial \overline{\Omega}} = \left(\frac{1 + \delta_{ij}}{2} \frac{\partial}{\partial \overline{\omega}_{ij}} \right).$$

Then

$$(2.10) \quad ds^2 = \sigma \left(Y^{-1} d\Omega Y^{-1} d\overline{\Omega} \right)$$

is a $Sp(g, \mathbb{R})$ -invariant metric on $\mathbb{H}_g$ (cf. [6]) and H. Maass [3] proved that its Laplacian is given by

$$(2.11) \quad \Delta = 4\sigma \left(Y^t \left(Y \frac{\partial}{\partial \overline{\Omega}} \right) \frac{\partial}{\partial \Omega} \right).$$

For $W = (w_{ij}) \in \mathbb{D}_g$, we write $dW = (dw_{ij})$ and $d\overline{W} = (d\overline{w}_{ij})$. We put

$$\frac{\partial}{\partial W} = \left(\frac{1 + \delta_{ij}}{2} \frac{\partial}{\partial w_{ij}} \right) \quad \text{and} \quad \frac{\partial}{\partial \overline{W}} = \left(\frac{1 + \delta_{ij}}{2} \frac{\partial}{\partial \overline{w}_{ij}} \right).$$

Using the Cayley transform $\Phi : \mathbb{D}_g \longrightarrow \mathbb{H}_g$, H. Maass proved (cf. [3]) that

$$(2.12) \quad ds_*^2 = 4\sigma \left((I_g - W\overline{W})^{-1} dW (I_g - \overline{W}W)^{-1} d\overline{W} \right)$$

is a G_* -invariant Riemannian metric on $\mathbb{D}_g$ and its Laplacian is given by

$$(2.13) \quad \Delta_* = \sigma \left((I_g - W\overline{W})^t \left((I_g - W\overline{W}) \frac{\partial}{\partial \overline{W}} \right) \frac{\partial}{\partial W} \right).$$

3. A Partial Cayley Transform

In this section, we present a partial Cayley transform of $\mathbb{D}_g \times \mathbb{C}^{(h,g)}$ onto $\mathbb{H}_g \times \mathbb{C}^{(h,g)}$ and prove that the action (1.3) of G^J on $\mathbb{H}_g \times \mathbb{C}^{(h,g)}$ is compatible with the *natural action* (cf. (3.5)) of the Jacobi group G_*^J on $\mathbb{D}_g \times \mathbb{C}^{(h,g)}$ via a partial Cayley transform.

From now on, for brevity we write $\mathbb{H}_{g,h} = \mathbb{H}_g \times \mathbb{C}^{(h,g)}$. We can identify an element $g = (M, (\lambda, \mu; \kappa))$ of G^J , $M = \begin{pmatrix} A & B \\ C & D \end{pmatrix} \in Sp(g, \mathbb{R})$ with the element

$$\begin{pmatrix} A & 0 & B & A^t \mu - B^t \lambda \\ \lambda & I_h & \mu & \kappa \\ C & 0 & D & C^t \mu - D^t \lambda \\ 0 & 0 & 0 & I_h \end{pmatrix}$$

of $Sp(g+h, \mathbb{R})$. This subgroup plays an important role in investigating the Fourier-Jacobi expansion of a Siegel modular form for $Sp(g+h, \mathbb{R})$ (cf. [4]) and studying the theory of Jacobi forms (cf. [1], [7-9], [15]).

We set

$$T_* = \frac{1}{\sqrt{2}} \begin{pmatrix} I_{g+h} & I_{g+h} \\ iI_{g+h} & -iI_{g+h} \end{pmatrix}.$$

We now consider the group G_*^J defined by

$$G_*^J = T_*^{-1} G^J T_*.$$

If $g = (M, (\lambda, \mu; \kappa)) \in G^J$ with $M = \begin{pmatrix} A & B \\ C & D \end{pmatrix} \in Sp(g, \mathbb{R})$, then $T_*^{-1} g T_*$ is given by

$$(3.1) \quad T_*^{-1} g T_* = \left(\begin{pmatrix} P_* & Q_* \\ \overline{Q}_* & \overline{P}_* \end{pmatrix}, \right),$$

where

$$P_* = \begin{pmatrix} P & \frac{1}{2} \{ Q^t (\lambda + i\mu) - P^t (\lambda - i\mu) \} \\ \frac{1}{2} (\lambda + i\mu) & I_h + i\frac{\kappa}{2} \end{pmatrix},$$

$$Q_* = \begin{pmatrix} Q & \frac{1}{2} \{ P^t (\lambda - i\mu) - Q^t (\lambda + i\mu) \} \\ \frac{1}{2} (\lambda - i\mu) & -i\frac{\kappa}{2} \end{pmatrix},$$

and P, Q are given by Formulas (2.4) and (2.5). From now on, we write

$$\left(\begin{pmatrix} P & Q \\ \overline{Q} & \overline{P} \end{pmatrix}, \left(\frac{1}{2} (\lambda + i\mu), \frac{1}{2} (\lambda - i\mu); -i\frac{\kappa}{2} \right) \right) = \left(\begin{pmatrix} P_* & Q_* \\ \overline{Q}_* & \overline{P}_* \end{pmatrix}, \right).$$

In other words, we have the relation

$$T_*^{-1} \left(\begin{pmatrix} A & B \\ C & D \end{pmatrix}, (\lambda, \mu; \kappa) \right) T_* = \left(\begin{pmatrix} P & Q \\ \overline{Q} & \overline{P} \end{pmatrix}, \left(\frac{1}{2} (\lambda + i\mu), \frac{1}{2} (\lambda - i\mu); -i\frac{\kappa}{2} \right) \right).$$

Let

$$H_{\mathbb{C}}^{(g,h)} = \left\{ (\xi, \eta; \zeta) \mid \xi, \eta \in \mathbb{C}^{(h,g)}, \zeta \in \mathbb{C}^{(h,h)}, \zeta + \eta^t \xi \text{ symmetric} \right\}$$

be the Heisenberg group endowed with the following multiplication

$$(\xi, \eta; \zeta) \circ (\xi', \eta'; \zeta') = (\xi + \xi', \eta + \eta'; \zeta + \zeta' + \xi^t \eta' - \eta^t \xi').$$

We define the semidirect product

$$SL(2g, \mathbb{C}) \ltimes H_{\mathbb{C}}^{(g,h)}$$

endowed with the following multiplication

$$\begin{aligned} & \left(\begin{pmatrix} P & Q \\ R & S \end{pmatrix}, (\xi, \eta; \zeta) \right) \cdot \left(\begin{pmatrix} P' & Q' \\ R' & S' \end{pmatrix}, (\xi', \eta'; \zeta') \right) \\ &= \left(\begin{pmatrix} P & Q \\ R & S \end{pmatrix} \begin{pmatrix} P' & Q' \\ R' & S' \end{pmatrix}, (\tilde{\xi} + \xi', \tilde{\eta} + \eta'; \zeta + \zeta' + \tilde{\xi}^t \eta' - \tilde{\eta}^t \xi') \right), \end{aligned}$$

where $\tilde{\xi} = \xi P' + \eta R'$ and $\tilde{\eta} = \xi Q' + \eta S'$.

If we identify $H_{\mathbb{R}}^{(g,h)}$ with the subgroup

$$\left\{ (\xi, \bar{\xi}; i\kappa) \mid \xi \in \mathbb{C}^{(h,g)}, \kappa \in \mathbb{R}^{(h,h)} \right\}$$

of $H_{\mathbb{C}}^{(g,h)}$, we have the following inclusion

$$G_*^J \subset SU(g, g) \ltimes H_{\mathbb{R}}^{(g,h)} \subset SL(2g, \mathbb{C}) \ltimes H_{\mathbb{C}}^{(g,h)}.$$

More precisely, if we recall $G_* = SU(g, g) \cap Sp(g, \mathbb{C})$ (cf. (2.6)), we see that the Jacobi group G_*^J is given by

$$(3.2) \quad G_*^J = \left\{ \left(\begin{pmatrix} P & Q \\ \bar{Q} & \bar{P} \end{pmatrix}, (\xi, \bar{\xi}; i\kappa) \right) \mid \begin{pmatrix} P & Q \\ \bar{Q} & \bar{P} \end{pmatrix} \in G_*, \xi \in \mathbb{C}^{(m,n)}, \kappa \in \mathbb{R}^{(m,m)} \right\}.$$

We define the mapping $\Theta : G^J \longrightarrow G_*^J$ by

$$(3.3) \quad \Theta \left(\begin{pmatrix} A & B \\ C & D \end{pmatrix}, (\lambda, \mu; \kappa) \right) = \left(\begin{pmatrix} P & Q \\ \bar{Q} & \bar{P} \end{pmatrix}, \left(\frac{1}{2}(\lambda + i\mu), \frac{1}{2}(\lambda - i\mu); -i\frac{\kappa}{2} \right) \right),$$

where P and Q are given by Formulas (2.4) and (2.5). We can see that if $g_1, g_2 \in G^J$, then $\Theta(g_1 g_2) = \Theta(g_1) \Theta(g_2)$.

According to [11, p. 250], G_*^J is of the Harish-Chandra type (cf. [5, p. 118]). Let

$$g_* = \left(\begin{pmatrix} P & Q \\ \bar{Q} & \bar{P} \end{pmatrix}, (\lambda, \mu; \kappa) \right)$$

be an element of G_*^J . Since the Harish-Chandra decomposition of an element $\begin{pmatrix} P & Q \\ R & S \end{pmatrix}$ in $SU(g, g)$ is given by

$$\begin{pmatrix} P & Q \\ R & S \end{pmatrix} = \begin{pmatrix} I_g & QS^{-1} \\ 0 & I_g \end{pmatrix} \begin{pmatrix} P - QS^{-1}R & 0 \\ 0 & S \end{pmatrix} \begin{pmatrix} I_g & 0 \\ S^{-1}R & I_g \end{pmatrix},$$

the P_*^+ -component of the following element

$$g_* \cdot \left(\begin{pmatrix} I_g & W \\ 0 & I_g \end{pmatrix}, (0, \eta; 0) \right), \quad W \in \mathbb{D}_g$$

of $SL(2g, \mathbb{C}) \ltimes H_{\mathbb{C}}^{(g,h)}$ is given by

$$(3.4) \quad \left(\begin{pmatrix} I_g & (PW + Q)(\bar{Q}W + \bar{P})^{-1} \\ 0 & I_g \end{pmatrix}, (0, (\eta + \lambda W + \mu)(\bar{Q}W + \bar{P})^{-1}; 0) \right).$$

We can identify $\mathbb{D}_g \times \mathbb{C}^{(h,g)}$ with the subset

$$\left\{ \left(\begin{pmatrix} I_g & W \\ 0 & I_g \end{pmatrix}, (0, \eta; 0) \right) \mid W \in \mathbb{D}_g, \eta \in \mathbb{C}^{(h,g)} \right\}$$

of the complexification of G_*^J . Indeed, $\mathbb{D}_g \times \mathbb{C}^{(h,g)}$ is embedded into P_*^+ given by

$$P_*^+ = \left\{ \left(\begin{pmatrix} I_g & W \\ 0 & I_g \end{pmatrix}, (0, \eta; 0) \right) \mid W = {}^t W \in \mathbb{C}^{(g,g)}, \eta \in \mathbb{C}^{(h,g)} \right\}.$$

This is a generalization of the Harish-Chandra embedding (cf. [5, p.119]). Hence G_*^J acts on $\mathbb{D}_g \times \mathbb{C}^{(h,g)}$ transitively by

$$(3.5) \quad \left(\begin{pmatrix} P & Q \\ \overline{Q} & \overline{P} \end{pmatrix}, (\lambda, \mu; \kappa) \right) \cdot (W, \eta) = \left((PW + Q)(\overline{Q}W + \overline{P})^{-1}, (\eta + \lambda W + \mu)(\overline{Q}W + \overline{P})^{-1} \right).$$

From now on, for brevity we write $\mathbb{D}_{g,h} = \mathbb{D}_g \times \mathbb{C}^{(h,g)}$. We define the map Φ_* of $\mathbb{D}_{g,h}$ into $\mathbb{H}_{g,h}$ by

$$(3.6) \quad \Phi_*(W, \eta) = \left(i(I_g + W)(I_g - W)^{-1}, 2i\eta(I_g - W)^{-1} \right), \quad (W, \eta) \in \mathbb{D}_{g,h}.$$

We can show that Φ_* is a biholomorphic map of $\mathbb{D}_{g,h}$ onto $\mathbb{H}_{g,h}$ which gives a partial bounded realization of $\mathbb{H}_{g,h}$ by the Siegel-Jacobi disk $\mathbb{D}_{g,h}$. We call this map Φ_* the *partial Cayley transform* of the Siegel-Jacobi disk $\mathbb{D}_{g,h}$.

Theorem 3.1. *The action (1.3) of G^J on $\mathbb{H}_{g,h}$ is compatible with the action (3.5) of G_*^J on $\mathbb{D}_{g,h}$ through the partial Cayley transform Φ_* . In other words, if $g_0 \in G^J$ and $(W, \eta) \in \mathbb{D}_{g,h}$,*

$$(3.7) \quad g_0 \cdot \Phi_*(W, \eta) = \Phi_*(g_* \cdot (W, \eta)),$$

where $g_* = T_*^{-1}g_0T_*$. We observe that Formula (3.7) generalizes Formula (2.9). The inverse of Φ_* is

$$(3.8) \quad \Phi_*^{-1}(\Omega, Z) = \left((\Omega - iI_g)(\Omega + iI_g)^{-1}, Z(\Omega + iI_g)^{-1} \right).$$

Proof. Let

$$g_0 = \left(\begin{pmatrix} A & B \\ C & D \end{pmatrix}, (\lambda, \mu; \kappa) \right)$$

be an element of G^J and let $g_* = T_*^{-1}g_0T_*$. Then

$$g_* = \left(\begin{pmatrix} P & Q \\ \overline{Q} & \overline{P} \end{pmatrix}, \left(\frac{1}{2}(\lambda + i\mu), \frac{1}{2}(\lambda - i\mu); -i\frac{\kappa}{2} \right) \right),$$

where P and Q are given by Formulas (2.4) and (2.5).

For brevity, we write

$$(\Omega, Z) = \Phi_*(W, \eta) \quad \text{and} \quad (\Omega_*, Z_*) = g_0 \cdot (\Omega, Z).$$

That is,

$$\Omega = i(I_g + W)(I_g - W)^{-1} \quad \text{and} \quad Z = 2i\eta(I_g - W)^{-1}.$$

Then we get

$$\begin{aligned}
\Omega_* &= (A\Omega + B)(C\Omega + D)^{-1} \\
&= \left\{ i A(I_g + W)(I_g - W)^{-1} + B \right\} \left\{ i C(I_g + W)(I_g - W)^{-1} + D \right\}^{-1} \\
&= \left\{ i A(I_g + W) + B(I_g - W) \right\} (I_g - W)^{-1} \\
&\quad \times \left[\left\{ i C(I_g + W) + D(I_g - W) \right\} (I_g - W)^{-1} \right]^{-1} \\
&= \left\{ (i A - B)W + (i A + B) \right\} \left\{ (i C - D)W + (i C + D) \right\}^{-1}
\end{aligned}$$

and

$$\begin{aligned}
Z_* &= (Z + \lambda\Omega + \mu)(C\Omega + D)^{-1} \\
&= \left\{ 2i\eta(I_g - W)^{-1} + i\lambda(I_g + W)(I_g - W)^{-1} + \mu \right\} \\
&\quad \times \left\{ i C(I_g + W)(I_g - W)^{-1} + D \right\}^{-1} \\
&= \left\{ 2i\eta + i\lambda(I_g + W) + \mu(I_g - W) \right\} (I_g - W)^{-1} \\
&\quad \times \left[\left\{ i C(I_g + W) + D(I_g - W) \right\} (I_g - W)^{-1} \right]^{-1} \\
&= \left\{ 2i\eta + (\lambda i - \mu)W + \lambda i + \mu \right\} \left\{ (i C - D)W + (i C + D) \right\}^{-1}.
\end{aligned}$$

On the other hand, we set

$$(W_*, \eta_*) = g_* \cdot (W, \eta) \quad \text{and} \quad (\widehat{\Omega}, \widehat{Z}) = \Phi_*(W_*, \eta_*).$$

Then

$$W_* = (PW + Q)(\overline{Q}W + \overline{P})^{-1} \quad \text{and} \quad \eta_* = (\eta + \lambda_* W + \mu_*)(\overline{Q}W + \overline{P})^{-1},$$

where $\lambda_* = \frac{1}{2}(\lambda + i\mu)$ and $\mu_* = \frac{1}{2}(\lambda - i\mu)$.

According to Formulas (2.4) and (2.5), we get

$$\begin{aligned}
\widehat{\Omega} &= i(I_g + W_*)(I_g - W_*)^{-1} \\
&= i \left\{ I_g + (PW + Q)(\overline{Q}W + \overline{P})^{-1} \right\} \left\{ I_g - (PW + Q)(\overline{Q}W + \overline{P})^{-1} \right\}^{-1} \\
&= i(\overline{Q}W + \overline{P} + PW + Q)(\overline{Q}W + \overline{P})^{-1} \\
&\quad \times \left\{ (\overline{Q}W + \overline{P} - PW - Q)(\overline{Q}W + \overline{P})^{-1} \right\}^{-1} \\
&= i \left\{ (P + \overline{Q})W + \overline{P} + Q \right\} \left\{ (\overline{Q} - P)W + \overline{P} - Q \right\}^{-1} \\
&= \left\{ (i A - B)W + (i A + B) \right\} \left\{ (i C - D)W + (i C + D) \right\}^{-1}
\end{aligned}$$

Therefore $\widehat{\Omega} = \Omega_*$. In fact, this result is the well-known fact (cf. Formula (2.9)) that the action (1.1) is compatible with the action (2.8) via the Cayley transform Φ . Here we gave

a proof of Formula (2.9) for convenience.

$$\begin{aligned}
\widehat{Z} &= 2i\eta_*(I_g - W_*)^{-1} \\
&= 2i(\eta + \lambda_*W + \mu_*)(\overline{Q}W + \overline{P})^{-1} \\
&\quad \times \left\{ I_g - (PW + Q)(\overline{Q}W + \overline{P})^{-1} \right\}^{-1} \\
&= 2i(\eta + \lambda_*W + \mu_*)(\overline{Q}W + \overline{P})^{-1} \\
&\quad \times \left\{ (\overline{Q}W + \overline{P} - PW - Q)(\overline{Q}W + \overline{P})^{-1} \right\}^{-1} \\
&= 2i(\eta + \lambda_*W + \mu_*) \left\{ (\overline{Q} - P)W + \overline{P} - Q \right\}^{-1}.
\end{aligned}$$

Using Formulas (2.4) and (2.5), we obtain

$$\widehat{Z} = \left\{ 2i\eta + (\lambda i - \mu)W + \lambda i + \mu \right\} \left\{ (iC - D)W + iC + D \right\}^{-1}.$$

Hence $\widehat{Z} = Z_*$. Consequently we get Formula (3.7). Formula (3.8) follows immediately from a direct computation. $\square$

For a coordinate $(\Omega, Z) \in \mathbb{H}_{g,h}$ with $\Omega = (\omega_{\mu\nu}) \in \mathbb{H}_g$ and $Z = (z_{kl}) \in \mathbb{C}^{(h,g)}$, we put

$$\begin{aligned}
\Omega &= X + iY, & X &= (x_{\mu\nu}), & Y &= (y_{\mu\nu}) \text{ real}, \\
Z &= U + iV, & U &= (u_{kl}), & V &= (v_{kl}) \text{ real}, \\
d\Omega &= (d\omega_{\mu\nu}), & dX &= (dx_{\mu\nu}), & dY &= (dy_{\mu\nu}), \\
dZ &= (dz_{kl}), & dU &= (du_{kl}), & dV &= (dv_{kl}), \\
d\overline{\Omega} &= (d\overline{\omega}_{\mu\nu}), & d\overline{Z} &= (d\overline{z}_{kl}),
\end{aligned}$$

$$\begin{aligned}
\frac{\partial}{\partial \Omega} &= \left(\frac{1 + \delta_{\mu\nu}}{2} \frac{\partial}{\partial \omega_{\mu\nu}} \right), & \frac{\partial}{\partial \overline{\Omega}} &= \left(\frac{1 + \delta_{\mu\nu}}{2} \frac{\partial}{\partial \overline{\omega}_{\mu\nu}} \right), \\
\frac{\partial}{\partial Z} &= \begin{pmatrix} \frac{\partial}{\partial z_{11}} & \cdots & \frac{\partial}{\partial z_{h1}} \\ \vdots & \ddots & \vdots \\ \frac{\partial}{\partial z_{1g}} & \cdots & \frac{\partial}{\partial z_{hg}} \end{pmatrix}, & \frac{\partial}{\partial \overline{Z}} &= \begin{pmatrix} \frac{\partial}{\partial \overline{z}_{11}} & \cdots & \frac{\partial}{\partial \overline{z}_{h1}} \\ \vdots & \ddots & \vdots \\ \frac{\partial}{\partial \overline{z}_{1g}} & \cdots & \frac{\partial}{\partial \overline{z}_{hg}} \end{pmatrix}.
\end{aligned}$$

Remark 3.1. The author proved in [13] that for any two positive real numbers A and B , the following metric

$$\begin{aligned}
(3.9) \quad ds_{g,h;A,B}^2 &= A \sigma \left(Y^{-1} d\Omega Y^{-1} d\overline{\Omega} \right) \\
&\quad + B \left\{ \sigma \left(Y^{-1} {}^t V V Y^{-1} d\Omega Y^{-1} d\overline{\Omega} \right) + \sigma \left(Y^{-1} {}^t (dZ) d\overline{Z} \right) \right. \\
&\quad \left. - \sigma \left(V Y^{-1} d\Omega Y^{-1} {}^t (d\overline{Z}) \right) - \sigma \left(V Y^{-1} d\overline{\Omega} Y^{-1} {}^t (dZ) \right) \right\}
\end{aligned}$$

is a Riemannian metric on $\mathbb{H}_{g,h}$ which is invariant under the action (1.3) of the Jacobi group G^J and its Laplacian is given by

$$(3.10) \quad \begin{aligned} \Delta_{n,m;A,B} = & \frac{4}{A} \left\{ \sigma \left(Y {}^t \left(Y \frac{\partial}{\partial \bar{\Omega}} \right) \frac{\partial}{\partial \Omega} \right) + \sigma \left(V Y^{-1} {}^t V {}^t \left(Y \frac{\partial}{\partial \bar{Z}} \right) \frac{\partial}{\partial Z} \right) \right. \\ & + \sigma \left(V {}^t \left(Y \frac{\partial}{\partial \bar{\Omega}} \right) \frac{\partial}{\partial Z} \right) + \sigma \left({}^t V {}^t \left(Y \frac{\partial}{\partial \bar{Z}} \right) \frac{\partial}{\partial \Omega} \right) \left. \right\} \\ & + \frac{4}{B} \sigma \left(Y \frac{\partial}{\partial Z} {}^t \left(\frac{\partial}{\partial \bar{Z}} \right) \right). \end{aligned}$$

We observe that Formulas (3.9) and (3.10) generalize Formulas (2.10) and (2.11). The following differential form

$$dv_{g,h} = (\det Y)^{-(g+h+1)} [dX] \wedge [dY] \wedge [dU] \wedge [dV]$$

is a G^J -invariant volume element on $\mathbb{H}_{g,h}$, where

$$[dX] = \wedge_{\mu \leq \nu} dx_{\mu\nu}, \quad [dY] = \wedge_{\mu \leq \nu} dy_{\mu\nu}, \quad [dU] = \wedge_{k,l} du_{kl} \quad \text{and} \quad [dV] = \wedge_{k,l} dv_{kl}.$$

Using the partial Cayley transform Φ_* and Theorem 3.1, we can find a G_*^J -invariant Riemannian metric on the Siegel-Jacobi disk $\mathbb{D}_{g,h}$ and its Laplacian explicitly which generalize Formulas (2.12) and (2.13). For more detail, we refer to [14].

4. The Canonical Automorphic Factors

The isotropy subgroup K_*^J at $(0,0)$ under the action (3.5) is

$$(4.1) \quad K_*^J = \left\{ \left(\begin{pmatrix} P & 0 \\ 0 & \bar{P} \end{pmatrix}, (0,0;\kappa) \right) \mid P \in U(g), \kappa \in \mathbb{R}^{(h,h)} \right\}.$$

The complexification of K_*^J is given by

$$(4.2) \quad K_{*,\mathbb{C}}^J = \left\{ \left(\begin{pmatrix} P & 0 \\ 0 & {}^t P^{-1} \end{pmatrix}, (0,0;\zeta) \right) \mid P \in GL(g, \mathbb{C}), \zeta \in \mathbb{C}^{(h,h)} \right\}.$$

By a complicated computation, we can show that if

$$(4.3) \quad g_* = \left(\begin{pmatrix} P & Q \\ \bar{Q} & \bar{P} \end{pmatrix}, (\lambda, \mu; \kappa) \right)$$

is an element of G_*^J , then the $K_{*,\mathbb{C}}^J$ -component of

$$g_* \cdot \left(\begin{pmatrix} I_g & W \\ 0 & I_g \end{pmatrix}, (0, \eta; 0) \right)$$

is given by

$$(4.4) \quad \left(\begin{pmatrix} P - (PW + Q)(\bar{Q}W + \bar{P})^{-1}\bar{Q} & 0 \\ 0 & \bar{Q}W + \bar{P} \end{pmatrix}, (0,0;\kappa_*) \right),$$

where

$$\begin{aligned} \kappa_* &= \kappa + \lambda {}^t \eta + (\eta + \lambda W + \mu) {}^t \lambda - (\eta + \lambda W + \mu) {}^t \bar{Q} ({}^t (\bar{Q}W + \bar{P})^{-1} {}^t (\eta + \lambda W + \mu)) \\ &= \kappa + \lambda {}^t \eta + (\eta + \lambda W + \mu) {}^t \lambda - (\eta + \lambda W + \mu) (\bar{Q}W + \bar{P})^{-1} \bar{Q} {}^t (\eta + \lambda W + \mu). \end{aligned}$$

Here we used the fact that $(\bar{Q}W + \bar{P})^{-1} \bar{Q}$ is symmetric.

For $g_* \in G_*^J$ given by (4.3) with $g_0 = \begin{pmatrix} P & Q \\ \overline{Q} & \overline{P} \end{pmatrix} \in G_*$ and $(W, \eta) \in \mathbb{D}_{g,h}$, we write

$$(4.5) \quad J(g_*, (W, \eta)) = a(g_*, (W, \eta)) b(g_0, W),$$

where

$$a(g_*, (W, \eta)) = (I_{2g}, (0, 0; \kappa_*)), \quad \text{where } \kappa_* \text{ is given in (4.4)}$$

and

$$b(g_0, W) = \left(\begin{pmatrix} P - (PW + Q)(\overline{Q}W + \overline{P})^{-1}\overline{Q} & 0 \\ 0 & \overline{Q}W + \overline{P} \end{pmatrix}, (0, 0; 0) \right).$$

Lemma 4.1. *Let*

$$\rho : GL(g, \mathbb{C}) \longrightarrow GL(V_\rho)$$

be a holomorphic representation of $GL(g, \mathbb{C})$ on a finite dimensional complex vector space V_ρ and $\chi : \mathbb{C}^{(h,h)} \longrightarrow \mathbb{C}^\times$ be a character of the additive group $\mathbb{C}^{(h,h)}$. Then the mapping

$$J_{\chi, \rho} : G_*^J \times \mathbb{D}_{g,h} \longrightarrow GL(V_\rho)$$

defined by

$$J_{\chi, \rho}(g_*, (W, \eta)) = \chi(a(g_*, (W, \eta))) \rho(b(g_0, W))$$

is an automorphic factor of G_^J with respect to χ and ρ .*

Proof. We observe that $a(g_*, (W, \eta))$ is a summand of automorphic, i.e.,

$$a(g_1 g_2, (W, \eta)) = a(g_1, g_2 \cdot (W, \eta)) + a(g_2, (W, \eta)),$$

where $g_1, g_2 \in G_*^J$ and $(W, \eta) \in \mathbb{D}_{g,h}$. Together with this fact, the proof follows from the fact that the mapping

$$J_\rho : G_* \times \mathbb{D}_g \longrightarrow GL(V_\rho)$$

defined by

$$J_\rho(g_0, W) := \rho(b(g_0, W)), \quad g_0 \in G_*, \quad W \in \mathbb{D}_g$$

is an automorphic factor of G_* . □

Example 4.1. Let $\mathcal{M}$ be a symmetric half-integral semi-positive definite matrix of degree h . Then the character

$$\chi_{\mathcal{M}} : \mathbb{C}^{(h,h)} \longrightarrow \mathbb{C}^\times$$

defined by

$$\chi_{\mathcal{M}}(c) = e^{-2\pi i \sigma(\mathcal{M}c)}, \quad c \in \mathbb{C}^{(h,h)}$$

provides the automorphic factor

$$J_{\mathcal{M}, \rho} : G_*^J \times \mathbb{D}_{g,h} \longrightarrow GL(V_\rho)$$

defined by

$$J_{\mathcal{M}, \rho}(g_*, (W, \eta)) = e^{-2\pi i \sigma(\mathcal{M}\kappa_*)} \rho(\overline{Q}W + \overline{P}),$$

where g_* is an element in G_*^J given by (4.3) and κ_* is given in (4.4). Using $J_{\mathcal{M}, \rho}$, we can define the notion of Jacobi forms on $\mathbb{D}_{g,h}$ of index $\mathcal{M}$ with respect to the Siegel modular group $T^{-1}Sp(g, \mathbb{Z})T$ (cf. [7-9]).

Remark 4.2. The P_*^- -component of

$$g_* \cdot \left(\begin{pmatrix} I_g & W \\ 0 & I_g \end{pmatrix}, (0, \eta; 0) \right)$$

is given by

$$(4.6) \quad \left(\left(\begin{pmatrix} I_g & 0 \\ (\overline{Q}W + \overline{P})^{-1}\overline{Q} & I_g \end{pmatrix}, \left(\lambda - (\eta + \lambda W + \mu)(\overline{Q}W + \overline{P})^{-1}\overline{Q}, 0; 0 \right) \right), \right.$$

where

$$P_*^- = \left\{ \left(\begin{pmatrix} I_g & 0 \\ W & I_g \end{pmatrix}, (\xi, 0; 0) \right) \mid W = {}^tW \in \mathbb{C}^{(g,g)}, \xi \in \mathbb{C}^{(h,g)} \right\}.$$

REFERENCES

- [1] M. Eichler and D. Zagier, *The Theory of Jacobi Forms*, Progress in Math., **55**, Birkhäuser, Boston, Basel and Stuttgart, 1985.
- [2] A. W. Knap, *Representation Theory of Semisimple Groups*, Princeton University Press, Princeton, New Jersey (1986).
- [3] H. Maass, *Die Differentialgleichungen in der Theorie der Siegelschen Modulformen*, Math. Ann. **126** (1953), 44–68.
- [4] I. Piatetski-Shapiro, *Automorphic Functions and the Geometry of Classical Domains*, Gordon-Breach, New York (1966).
- [5] I. Satake, *Algebraic Structures of Symmetric Domains*, Kano Memorial Lectures 4, Iwanami Shoton, Publishers and Princeton University Press (1980).
- [6] C. L. Siegel, *Symplectic Geometry*, Amer. J. Math. **65** (1943), 1-86; Academic Press, New York and London (1964); *Gesammelte Abhandlungen*, no. 41, vol. II, Springer-Verlag (1966), 274-359.
- [7] J.-H. Yang, *The Siegel-Jacobi Operator*, Abh. Math. Sem. Univ. Hamburg **63** (1993), 135–146.
- [8] J.-H. Yang, *Singular Jacobi Forms*, Trans. Amer. Math. Soc. **347** (6) (1995), 2041–2049.
- [9] J.-H. Yang, *Construction of vector valued modular forms from Jacobi forms*, Canadian J. of Math. **47** (6) (1995), 1329–1339.
- [10] J.-H. Yang, *A geometrical theory of Jacobi forms of higher degree*, Proceedings of Symposium on Hodge Theory and Algebraic Geometry (edited by Tadao Oda), Sendai, Japan (1996), 125–147 or Kyungpook Math. J. **40** (2) (2000), 209–237 or arXiv:math.NT/0602267.
- [11] J.-H. Yang, *The Method of Orbits for Real Lie Groups*, Kyungpook Math. J. **42** (2) (2002), 199–272 or arXiv:math.RT/0602056.
- [12] J.-H. Yang, *A note on a fundamental domain for Siegel-Jacobi space*, Houston Journal of Mathematics, **32** (3) (2006), 701–712.
- [13] J.-H. Yang, *Invariant metrics and Laplacians on Siegel-Jacobi space*, arXiv:math.NT/0507215 v1 or revised version (2006).
- [14] J.-H. Yang, *Invariant metrics and Laplacians on Siegel-Jacobi disk*, arXiv:math.NT/0507217 v1 or revised version (2006).
- [15] C. Ziegler, *Jacobi Forms of Higher Degree*, Abh. Math. Sem. Hamburg **59** (1989), 191–224.

DEPARTMENT OF MATHEMATICS, INHA UNIVERSITY, INCHEON 402-751, KOREA

E-mail address: jhyang@inha.ac.kr